

GALWAYELLA, A NEW FORAMINIFERAL GENUS AND NEW NAMES FOR TWO FORAMINIFERAL HOMONYMS

R. TIMOTHY PATTERSON AND RANI H. PETTIS

Department of Earth and Space Sciences, University of California,

Los Angeles, CA 90024

ABSTRACT

Galwayella, a new genus of the Oolininae differs from other genera of this family by its distinct trigonal cross-section. The type species of the new genus is *Lagena trigonoelliptica* Balkwill and Millett, 1884.

New names are proposed for two homonyms in the Oolininae and Nodosariinae: *Lagenosolenia incomposita* for *Lagena marginata* (Montagu) var. *spinifera* Earland, and *Dentalina ariena* for *Dentalina intorta* (Dervieux).

INTRODUCTION

The authors are currently studying benthic foraminifera from the Rio Grande Rise in the southwest Atlantic Ocean, and from the Benham Rise in the western Philippine Basin of the western Pacific Ocean. While identifying Quaternary species of the Oolininae and Nodosariinae a new genus has been differentiated and some specific taxa were found to be homonyms of previously described species. We are herein describing the new genus and proposing replacement names for the junior homonyms.

SYSTEMATIC DESCRIPTION

Suborder ROTALINA Delage and Herouard, 1896
Superfamily NODOSARIACEA Ehrenberg, 1838

Family GLANDULINIDAE Reuss, 1860

Subfamily OOLININAE Loeblich and Tappan, 1961
Galwayella, n. gen.

Trigonulina SEGUENZA, 1862, p. 74 (not d'Orbigny, 1846).

Type species. Lagena trigonoelliptica Balkwill and Millett, 1884, p. 81, 87, pl. 3, fig. 8 (as *L. marginata* var. *trigono-elliptica* on p. 87).

Diagnosis. An oolinid with a distinct trigonal cross-section.

Description. Test free, unilocular, triangular in section, apices mayor may not be keeled, subfusiform longitudinally, broadest centrally and tapering at ends; wall calcareous, hyaline, smooth; aperture a curved narrow slit to triangular, apertural region may be compressed, with a short centrally located entosolenian tube; neck mayor may not be distinct.

Derivation of name. From Galway Bay, Eire, plus *-ella*, diminutive, with reference to the type locality of the type species of the microfossil.

Remarks. *Galwayella*, n. genus, differs from *Fissurina* Reuss, 1850 in the distinct trigonal cross-section and the trigonal aperture of many species. *Trigonulina* Seguenza (1862), with type species *Trigonulina oblonga* Seguenza (1862) as designated by Cushman (1928, p. 193), may be congeneric. Unfortunately, *Trigono-*

lina Seguenza is a homonym of *Trigonulina* d'Orbigny (1846, Mollusca) and is therefore invalid. Seguenza (1862) lists no depository for *Trigonulina oblonga*. As no specimens of *T. oblonga* were available for this study, its placement in *Galwayella*, although probable, is not certain. *Lagena trigonoelliptica* Balkwill and Millett (1884), was available for study from the Quaternary of DSDP Site 357 in the southwest Atlantic, and was therefore chosen as the type species of *Galwayella* (Figs. 1-4).

Certain species formerly included in *Trigonulina* Seguenza and *Lagena* Walker and Jacob (in Kanmacher, 1798) are herein transferred to *Galwayella*. These include *Trigonulina biangulata* Seguenza, now *Galwayella biangulata* (Seguenza), *Trigonulina globosa* Seguenza, now *G. globosa* (Seguenza), *Trigonulina oblonga* Seguenza, now *G. oblonga* (Seguenza), *Trigonulina regularis* Seguenza, now *G. regularis* (Seguenza), *Lagena trigona* Koch, now *G. trigona* (Koch), *Lagena trigonobicarinata* Balkwill and Millett, now *G. trigonobicarinata* (Balkwill and Millett), *Lagena trigonoorbignyana* Balkwill and Millett, now *G. trigonoorbignyana* (Balkwill and Millett), *Lagena trigonoornata* Brady, now *G. trigonoornata* (Brady) and *Lagena trigonopulchella* Balkwill and Millett, now *G. trigonopulchella* (Balkwill and Millett).

Lagenosolenia McCulloch, 1977

Lagenosolenia incomposita, n. name

Lagenosolenia incomposita new name for *Lagena marginata* (Montagu) var. *spinifera* EARLAND, 1934, p. 156, pl. 7, figs. 3, 4, not *Lagena aspera* Reuss var. *spinifera* CHAPMAN 1895, p. 27, pl. 1, fig. 4.

Derivation of name. From the Latin, *incompositus*, disarranged, confused, with reference to the muddled nomenclatorial history of the species.

Remarks. *Lagena marginata* (Montagu) var. *spinifera* Earland is a primary homonym of *Lagena aspera* Reuss var. *spinifera* Chapman. Although herein transferred to *Lagenosolenia*, the species *spinifera* was a homonym when coined by Earland in 1934 and is therefore unavailable. The new name *Lagenosolenia incomposita* is proposed as a replacement.


Family NODOSARIIDAE Ehrenberg, 1838

Subfamily NODOSARIINAE Ehrenberg, 1838

Dentalina Risso, 1826

Dentalina ariena, n. name

Dentalina ariena new name for *Dentalina intorta* (Dervieux) BARKER, 1960, (ex *Nodosaria intorta* Dervieux, 1894) p. 130, pl. 62, figs. 27-31, not *Dentalina intorta* TERQUEM, 1870, p. 364, pl. 27, figs. 26-34.



FIGURES 1-4. *Galwayella trigonoelliptica* (Balkwill and Millett). 1 Side view of hypotype (USNM 383281); x 300. 2 Apertural view of same specimen showing trigonal opening; from DSDP Site 357, Core 1, Section 2, 80-86 cm; x 300. 3,4 Side and apertural view of specimen figured by Balkwill and Millett (1884, pl. 3, fig. 8); from the shoresands of Galway Bay, near Salthill, about two miles west of Galway, Eire; x 145.

Derivation of name. From the Latin, *ariena*, fruit of pala, plantain, with reference to the shape of the test.

ACKNOWLEDGMENTS

This research was supported by a Canadian Natural Sciences and Engineering Research Council Postgraduate Scholarship to R. T. Patterson. Samples from the Deep Sea Drilling Project were supplied through the assistance of the National Science Foundation. National Science Foundation Grant EAR-8306170 to Alfred R. Loeblich, Jr. and Helen Tappan provided support. We would also like to thank Helen Tappan, Alfred R. Loeblich, Jr., and Stephen Culver for critically reviewing the manuscript.

REFERENCES

- BALKWILL, F. P., and MILLETT, F. W., 1884, The foraminifera of Galway: Part 2; *Journal of Microscopy and Natural Science*, v. 3, p. 78-90.
- BARKER, R. W., 1960, Taxonomic notes on the species figured by H. B. Brady in his Report on the Foraminifera dredged by H.M.S. *Challenger* during the years 1873-1876; *Society of Economic Paleontologists and Mineralogists Special Publication No. 9*, Tulsa, Oklahoma, 238 p.
- BRADY, H. B., 1884, Report on the Foraminifera dredged by H.M.S. *Challenger* during the years 1873-1876; Reports of the Scientific Results of the Voyage of the H.M.S. *Challenger*, v. 9, (Zoology), 814 p.
- CHAPMAN, F., 1895, On some foraminifera obtained by the Royal Indian Marine Survey's S.S. "Investigator," from the Arabian Sea, near the Laccadive Islands: Proceedings of the Zoological Society of London, p. 4-55.
- CUSHMAN, J. A., 1928, Foraminifera their classification and economic use: Cushman Laboratory For Foraminiferal Research, Special Publication No. 1, 401 p.
- EARLAND, A., 1934, Foraminifera; Part 3- The Falklands sector of the Antarctic (excluding South Georgia); *Discovery Reports*, v. 10, p. 1-208.
- SEGUENZAG., 1862, Dei terreni Terziarii del distretto di Messina; Parte II - Descrizione dei foraminiferi monotalmici delle marne mioceniche del distretto di Messina: T. Capra, Messina, Italia, 84 p.
- TERQUEM, O., 1870, Troisieme memoire sur les foraminiferes du systeme oolithique, comprenant les genres *Frondicularia*, *Flabellina*, *Nodosaria*, *Dentalina*, etc. de la zone a *Ammonites parkinsoni* de Fontoy (Moselle): Academie Imperiale de Metz, Memoires, annee 51 (ser. 2, annee 18) 1869-1870, p. 299-380.
- WALKER, G., and JACOB, E., in KANMACHER, 1798, Adam's essays on the microscope: Dillon and Keaton, London, England, 2nd edition, 712 p.

Received October 18, 1984
Accepted November 19, 1984